

Central Forces

$$\vec{F} = m_{\text{eff}} \vec{a} = m_{\text{eff}} \left[(\ddot{r} - r\dot{\theta}^2) \hat{r} + (r\ddot{\theta} + 2\dot{r}\dot{\theta}) \hat{\theta} \right].$$

Central force


$$\vec{F} = \vec{F}(\vec{r}) = \left(\frac{k}{r^2} \right) \hat{r}.$$

$$\Rightarrow m_{\text{eff}} \ddot{r} - m_{\text{eff}} r \dot{\theta}^2 = \frac{k}{r^2}. \quad \text{--- (i)}$$

$$\& m_{\text{eff}} r \ddot{\theta} + 2m_{\text{eff}} \dot{r} \dot{\theta} = 0. \quad \text{--- (ii)}$$

$$\text{Eq. (ii)} \Rightarrow \frac{d}{dt} (m_{\text{eff}} r^2 \dot{\theta}) = 0.$$

$$\therefore m_{\text{eff}} r^2 \dot{\theta} = \text{const} = L \text{ (ang. mom.)}.$$



Area swept by the radius vector in time dt

$$\text{is } dA = \frac{1}{2} r \cdot r d\theta = \frac{r^2 d\theta}{2}$$

$$\therefore \text{Areal velocity} = \frac{1}{2} r^2 \dot{\theta} = \frac{L}{2m_{\text{eff}}} = \text{constant}.$$

Also,

$$\begin{aligned}
 E &= T + V \\
 &= \frac{1}{2} m_{\text{eff}} \dot{r}^2 + \frac{1}{2} m_{\text{eff}} r^2 \dot{\theta}^2 + V(r) \\
 &= \frac{1}{2} m_{\text{eff}} \dot{r}^2 + \frac{L^2}{2m_{\text{eff}} r^2} + V(r)
 \end{aligned}$$

We had from (i),

$$m_{\text{eff}} \ddot{r} = - \frac{d}{dr} \left(V + \frac{L^2}{2m_{\text{eff}} r^2} \right)$$

Multiply both sides by \dot{r}

$$\& \text{ use } \dot{r} \frac{d}{dr} = \frac{d}{dt}$$

$$\Rightarrow m_{\text{eff}} \dot{r} \ddot{r} = - \frac{d}{dt} \left(V + \frac{L^2}{2m_{\text{eff}} r^2} \right)$$

$$\text{or, } \frac{d}{dt} \left[\frac{1}{2} m_{\text{eff}} \dot{r}^2 + \frac{L^2}{2m_{\text{eff}} r^2} + V \right] = 0$$

$$\therefore \frac{1}{2} m_{\text{eff}} \dot{r}^2 + \frac{L^2}{2m_{\text{eff}} r^2} + V = \text{const.}$$

$$E = \text{const.}$$

$$\Rightarrow \int_{r_0}^r \frac{dr}{\left[\frac{2}{m_{\text{eff}}} \left(E - V - \frac{L^2}{2m_{\text{eff}} r^2} \right) \right]^{1/2}} = \int_0^t dt = t$$

Also, $\dot{\theta} = \frac{L}{m_{\text{eff}} r^2}$

$$\Rightarrow \int_{\theta_0}^{\theta} d\theta = \int_0^t \frac{L}{m_{\text{eff}} r^2} dt$$

Also, $d\theta = \frac{d\theta}{dt} \frac{dt}{dr} dr = \frac{\dot{\theta}}{\dot{r}} dr$

$$= \frac{L}{m_{\text{eff}} r^2 \dot{r}} dr$$

$$\therefore d\theta = \frac{(L/r^2)}{m_{\text{eff}} \dot{r}} dr$$


But $m_{\text{eff}} \dot{r} = \left[2m_{\text{eff}} \left(E - V - \frac{L^2}{2m_{\text{eff}} r^2} \right) \right]^{1/2}$

$$\Rightarrow \int_{\theta_0}^{\theta} d\theta = \int_{r_0}^r \frac{L/r^2}{\left[2m_{\text{eff}} \left(E - V - \frac{L^2}{2m_{\text{eff}} r^2} \right) \right]^{1/2}} dr$$

$$F(r) = \frac{k}{r^2}$$

$$\therefore V(r) = + \frac{k}{r}$$

$$V_e = \frac{k}{r} + \frac{L^2}{2m_e r^2}$$



$$m_{\text{eff}} \ddot{r} = F(r) + \frac{L^2}{m_{\text{eff}} r^3} \quad \text{--- (A)}$$

$$\text{Let } u = \frac{1}{r}$$

$$\frac{du}{d\theta} = -\frac{1}{r^2} \frac{dr}{d\theta}$$

$$= -\frac{1}{r^2} \frac{dr}{dt} \frac{dt}{d\theta}$$

$$\therefore \frac{du}{d\theta} = -\frac{\dot{r}}{r^2 \dot{\theta}} = -\frac{m_{\text{eff}}}{L} \dot{r}$$

$$\frac{d^2 u}{d\theta^2} = \frac{d}{d\theta} \left(\frac{du}{d\theta} \right)$$

$$= \frac{d}{d\theta} \left(-\frac{m_{\text{eff}}}{L} \dot{r} \right)$$

$$= \frac{d}{dt} \left(-\frac{m_{\text{eff}}}{L} \dot{r} \right) \frac{dt}{d\theta}$$

$$= -\frac{m_{\text{eff}}}{L} \ddot{r}$$

$$= -\frac{m_{\text{eff}} r^2}{L^2} \ddot{r}$$

$$\therefore \frac{d^2 u}{d\theta^2} + u = -\frac{m_{\text{eff}}}{L^2 u^2} F\left(\frac{1}{u}\right)$$

$$F(r) = \frac{|k|}{r^2}$$

$$F\left(\frac{1}{u}\right) = |k|u^2$$

$$\therefore \frac{d^2u}{d\theta^2} + u = \frac{|k|m_{\text{eff}}}{L^2}$$

$$\text{Let } y = u - \frac{|k|m_{\text{eff}}}{L^2}$$

$$\Rightarrow \frac{d^2y}{d\theta^2} + y = 0$$

$$y = A \cos(\theta - \theta_0)$$

$$\therefore \frac{1}{r} = \frac{|k|m_{\text{eff}}}{L^2} + A \cos(\theta - \theta_0)$$

$$\text{or, } \frac{L^2/|k|m_{\text{eff}}}{r} = 1 + \frac{L^2 A \cos(\theta - \theta_0)}{|k|m_{\text{eff}}}$$

$$\frac{1}{r} = 1 + e \cos \theta$$

$$l = \frac{L^2}{|k|m} \quad ; \quad e = \frac{L^2 A}{|k|m}$$

Extremal values of $\cos\theta = \pm 1$. (7)

L14

$$\left. \begin{aligned} \frac{1}{r_1} &= \frac{m|k|}{L^2} + A, \\ \frac{1}{r_2} &= \frac{m|k|}{L^2} - A. \end{aligned} \right\} \text{turning points.}$$

Turning points are also roots of the equation,

$$E - V_e(\vec{r}) = E + \frac{|k|}{r} - \frac{L^2}{2mr^2} = 0.$$

The roots are,

$$\frac{1}{r_{1,2}} = \frac{m|k|}{L^2} \pm \frac{m|k|}{L^2} \sqrt{1 + \frac{2EL^2}{mk^2}}.$$

$$\therefore \frac{1}{r_1} - \frac{1}{r_2} = 2A = \frac{2m|k|}{L^2} \sqrt{1 + \frac{2EL^2}{mk^2}}.$$

$$\therefore A = \frac{m|k|}{L^2} \sqrt{1 + \frac{2EL^2}{mk^2}}.$$

\Rightarrow Eccentricity, e

$$\begin{aligned} &= \frac{L^2 A}{m|k|} = \frac{L^2}{m|k|} \frac{m|k|}{L^2} \sqrt{1 + \frac{2EL^2}{mk^2}} \\ &= \sqrt{1 + \frac{2EL^2}{mk^2}}. \end{aligned}$$

(1)

(2)

AJ

$E = 0$ Circle $E = -\frac{mk^2}{2L^2}$

$E = 1$ Parabola $E = 0$

$E < 1$ Ellipse $E < 0, \neq -\frac{mk^2}{2L^2}$

$E > 1$ Hyperbola $E > 0$